


0 : 59 : 53




Question 1 / 10

In the figure shown the inclined plane is frictionless. If $\theta = 45^\circ$, $m_2 = 3 \text{ kg}$, $m_1 = 6 \text{ kg}$. The tension (in N) in the cord is: (Use $g = 9.8 \text{ m/s}^2$)

1. 38.5
2. 30.8
3. 33.5
4. 35.7
5. 37.4

Next




Question 2 / 10

Three objects A, B and C attached to each other and have equal masses $m=1$ kg. If the objects are pulled vertically upward with force $F=3$ N. The acceleration (in m/s^2) of object C is:

1. 1
2. 2
3. 3
4. 4
5. 5

00 : 58 : 11




Question 3 / 10

If the coefficient of static friction between the block shown and the wall is 0.1 and the mass of the block is 0.6 Kg. Find the minimum value for the force F (in N) needed to prevent the block from motion. (Use $g = 9.8 \text{ m/s}^2$)

1. 19.6
2. 29.4
3. 39.2
4. 58.8
5. 68.6

Previous

Next



4/10

Question:

For the Atwood machine shown $m_1=1$ kg $m_2=3$ kg. The tension (in N) in the string is: (Use $g=9.8$ m/s²)

1. 13.1
2. 14.7
3. 15.7
4. 16.3
5. 16.8

Previous

Next

00 : 58 : 27



5/10

Question

In the figure shown if the angle between the acceleration ($a = 14 \text{ m/s}^2$) and the radius ($R = 44 \text{ m}$) is ($\theta = 60^\circ$). Find the speed (in m/s) of the particle shown.

1. ≈ 14.96
2. ≈ 16.31
3. ≈ 17.55
4. ≈ 18.71
5. ≈ 20.83

Next

Start

Calculator

<http://www2.bas.edu>

acer

00 : 57 : 52

Question 6 / 10

A block with mass M is whirled on the end of a thin rigid rod that moves at a constant speed in a vertical circle with radius 8 m . At the top of the circle, the tension in the rod is twice the weight of the block. What is the speed (in m/s) of the block? (Use $g = 9.8\text{ m/s}^2$)

- 1. 13.3
- 2. 14.3
- 3. 15.3
- 4. 16.3
- 5. 20.3

Previous Next

Submit Answer

00 : 48 : 0

Question 7 / 10

A 4.0-kg object rests on the floor of an elevator which is accelerating downward at rate of 1.8 m/s^2 . What is the magnitude of the force (in N) the object exerts on the elevator? (Use $g=9.8 \text{ m/s}^2$)


- 1. 16
- 2. 24
- 3. 32
- 4. 40
- 5. 48

Previous

Next

إضافة ملاحظة

00 : 47 : 47




Question 8 / 10

A block is pushed across a horizontal surface by the force shown. If the coefficient of kinetic friction between the block and the surface is 0.10, $F = 20\text{ N}$, $\theta = 30^\circ$, and $M = 3.0\text{ kg}$. The magnitude of the acceleration (in m/s^2) of the block is: (Use $g = 9.8\text{ m/s}^2$)

1. 4.5
2. 3.8
3. 3.2
4. 2.5
5. 1.8

Previous

Next



Question 9 / 10


For the conical pendulum shown if $m=80$ kg, $\theta=40^\circ$ and its length is 10 m. The speed (in m/s) of m is: (Use $g=9.8$ m/s²)

1. 6.3
2. 6.8
3. 7.3
4. 11.6
5. 16.9

Question 10 / 10

The cylinder shown is rotating around its axis (the dashed line). If the cylinder is opened from the bottom find the minimum velocity needed for the cylinder to have such that the mass does not fall down. (Radius of cylinder $R=4$ m, static friction coefficient between block and cylinder wall is $\mu_s=0.98$, $M=4$ kg). (Use $g=9.8$ m/s²)

- 1. 3.2
- 2. 4.5
- 3. 5.5
- 4. 6.3
- 5. 7.1



Previous

إرسال ملاحظتك

Q1

$m_1 = 6\text{ kg}, m_2 = 3\text{ kg}, \theta = 45$

$g = 9.8\text{ m/s}^2, T = ??$

$w_1 = 58.8\text{ N}$

$w_2 = 29.4\text{ N}$

$\sum F_{yD} = m_1 a$

$w_1 - T = m_1 a$

$58.8 - T = 6a \dots \textcircled{1}$

$\sum F_{xII} = m_2 a$

$T - w_2 \sin \theta = m_2 a$

$T - 29.4 \sin 45 = 3a$

$T - 20.8 = 3a \dots \textcircled{2}$

⊕ ⊕ ⊕ ⊕ ⊕

$\Rightarrow 38 = 4a \rightarrow a = 4.2\text{ m/s}^2$


⊕ ⊕ ⊕ ⊕ ⊕

$58.8 - T = 6(4.2) \rightarrow T = 33.5\text{ N}$

Q2 $m_a = m_b = m_c = 1\text{ kg}, F = 3\text{ N}$

$F = ma$

$3 = 3a \rightarrow a = 1\text{ m/s}^2$



! عذرا، اس کا جواب نہیں

Q3 $m = 0.6 \text{ kg}$, $\mu_s = 0.1$, $g = 9.8 \text{ m/s}^2$

$$\sum F_y = 0$$

$$f_s - w = 0 \rightarrow f_s = w = 5.9 \text{ N}$$


$$\sum F_x = 0$$

~~$$F - w = 0$$~~

$$F - n = 0 \rightarrow F = n = 58.8 \text{ N}$$

$$f_s = \mu_s n$$

$$5.9 = 0.1 n \rightarrow n = 59 \text{ N}$$



Q4 $M_1 = 1 \text{ kg}$, $M_2 = 3 \text{ kg}$, $g = 9.8 \text{ m/s}^2$

$$w_1 = 9.8 \text{ N}$$

$$w_2 = 29.4 \text{ N}$$

$$w_2 - T = m_2 a$$


$$29.4 - T = 3a \dots \textcircled{1}$$

~~$$T - w_1 = m_1 a$$~~

$$T - 9.8 = a \dots \textcircled{2}$$

$$19.6 = 4a \rightarrow a = 4.9 \text{ m/s}^2$$


$$29.4 - T = 14.7 \rightarrow T = 14.7 \text{ N}$$



Q5 $a_{\text{tot}} = 14 \text{ m/s}^2$, $R = 44 \text{ m}$, $\theta = 60^\circ$
Find N ?

$$a_r = a_{\text{tot}} \cos \theta$$

$$= 14 \cos 60 \rightarrow a_r = 7 \text{ m/s}^2$$




$$a_r = \frac{v^2}{r} \rightarrow 7 = \frac{v^2}{44} \rightarrow v = 17.55 \text{ m/s}$$

Q6 $R = 8 \text{ m}$, $g = 9.8 \text{ m/s}^2$

$$T = 2Mg$$

$$\sum F_r = m a_r$$



$$T + W = M \frac{v^2}{r}$$

$$2Mg + Mg = M \frac{v^2}{r}$$

$$3Mg = M \frac{v^2}{r}$$

$$29.4 = \frac{v^2}{8} \rightarrow v = 15.34 \text{ m/s}$$


Q7 $m = 4 \text{ kg}$, $\vec{a} = 1.8 \text{ m/s}^2$, $g = 9.8$

$$\sum F_y = ma$$

$$W - n = ma$$

$$39.2 - n = 7.2$$

$$n = 32 \text{ N}$$



Q8 $m = 3 \text{ kg}$, $F = 20 \text{ N}$, $\theta = 30^\circ$, $\mu_k = 0.1$
 $g = 9.8 \text{ m/s}^2$


$$\sum F_y = 0$$

$$n - W - F \sin 30 = 0$$

$$n - 29.4 - 10 = 0$$

$$n = 39.4 \text{ N}$$

$$\left. \begin{aligned} f_k &= \mu_k n \\ &= 0.1 \times 39.4 \\ f_k &= 3.9 \text{ N} \end{aligned} \right\}$$



$$\sum F_x = ma$$

$$F \cos 30 - f_k = ma$$


$$17.3 - 3.9 = 3a \rightarrow a = 4.5 \text{ m/s}^2$$

Q9) $m = 80 \text{ kg}$, $\theta = 40^\circ$, $L = 10 \text{ m}$, $g = 9.8 \text{ m/s}^2$

$$\sum F_y = 0$$

$$T \cos 40^\circ = W = 784$$

$$T \cos 40^\circ = 784 \Rightarrow T = 1023.4 \text{ N}$$



$$\sum F_r = m a_r$$

$$T \sin 40^\circ = m \frac{v^2}{r}$$

$$10 \times 23.4 \sin 40^\circ = 80 \frac{v^2}{6.4}$$

$$\sin 40^\circ = \frac{r}{L}$$

$$r = 6.4 \text{ m}$$

$$\Rightarrow v = 7.3 \text{ m/s}$$

Q10) $R = 4 \text{ m}$, $\mu_s = 0.8$, $M = 4 \text{ kg}$, $g = 9.8 \text{ m/s}^2$

$$\sum F_y = 0$$

$$f_s - W = 0 \Rightarrow f_s = W = 39.2 \text{ N}$$


$$\sum F_r = m \frac{v^2}{r}$$

$$40 = 4 \frac{v^2}{4}$$

$$f_s = \mu_s N$$

$$39.2 = 0.8 N$$

$$N = 49 \text{ N}$$



$$v_{\min} = 6.3 \text{ m/s}$$

اعداد بظاہر - کلاس، انمولین